Coastal AgroBusiness, Inc.

Product Label

N-Put

Active Ingredients:

Pseudomonas siliginis **.....1x10^6 CFU/mL**

Rhizobium tropici bv populus **..... 1x10^6 CFU/mL**

Inert Ingredients:

Aqueous Carrier 99.0%

Weight per gallon: 8.36 lbs.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Read the entire label for additional precautionary statements.

SHAKE WELL BEFORE USING

Manufactured By:
Coastal AgroBusiness, Inc.
Greenville, North Carolina

GENERAL INFORMATION

N-Put is a ready-to-use blend of 2 biologicals that live inside the plant. The biologicals convert atmospheric nitrogen to a usable form inside the plant and increase the metabolism of phosphorous. **N-Put** increases nitrogen availability to feed your crop all season long. **N-Put** is not a fertilizer substitute.

PRECAUTIONARY STATEMENTS / HAZARDS TO HUMANS

CAUTION. Avoid contact with eyes, skin, or clothing. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse. Before using this product, read all precautions, directions for use, conditions of sale - limited warranty, and limitations of liability and remedies.

FIRST AID

If swallowed: Call a physician immediately. Only induce vomiting at the instruction of a physician. Never give anything by mouth to an unconscious person.

If in eyes: Flush eyes with copious amounts of clean water for at least 15 minutes. Remove contacts if present, and easy to do. Do not let the victim rub their eye. Get medical attention.

If on skin: Remove contaminated clothing and wash skin with soap and water. If irritation develops, get medical attention.

If inhaled: Remove the person from the area of exposure. Give artificial respiration if necessary. If breathing is difficult, give oxygen if a qualified operator is available. Seek medical attention.

STORAGE AND DISPOSAL

Store in original package only. Only open the package when ready to use. Store in a cool, dry location out of direct sunlight in temperatures between 35 °F and 70°F in the original packaging. Do not contaminate water, food, or feed by storage or disposal. Do not contaminate water by runoff from equipment wash waters or spray materials. Wastes resulting from this product may be disposed of on-site or at an approved waste disposal facility.

Container disposal: Empty packaging should be offered for recycling or disposal in accordance with federal, state, and local regulations. If packaging is burned, stay out of smoke.

AGRICULTURAL USE REQUIREMENTS AND USER SAFETY RECOMMENDATIONS

Use of waterproof gloves and protective eyewear is recommended when handling this product.

DIRECTIONS FOR USE

Shake or agitate contents of package to mix thoroughly. Open package only when ready to use. Add to tank last. Apply product foliarly with 10 - 20 gallons of water. Apply product in-furrow with the seed in enough liquid carrier to ensure good seed contact.

Application rate: 16 oz. per acre foliarly or in-furrow. For best results, apply foliarly early in the growing cycle with enough foliage to capture the application. See the Rate Chart below for further application frequency and timing notes.

Crop	Total Annual Application	Application Frequency and Timing
Wheat, Barley, Corn, Oats, Rye, Triticale, Canola, Soybeans, Flax, Rapeseed, Lentils, Peas, Chickpeas, Kidney beans, Sunflowers, Buckwheat	16 to 32 fl. oz./acre	Apply 1 - 2 times between 3-leaf stage until pre-senescence
Fruiting Vegetables (Tomato, Pepper) & Cucurbit Vegetables (Cucumber, Pumpkin, Melons, Squash)	16 to 64 fl. oz./acre	Apply 2 - 4 times between 3-leaf stage until pre-senescence. Apply during active growth.
Small Fruits and Berries (Cranberries, Blueberries, Raspberries, Strawberries)	16 to 64 fl. oz./acre	Apply 2 - 8 times during active growth.
Apples, Pears, Peaches, Plums, Cherries, Table grapes, Wine grapes, Millet, Sorghum, Alfalfa, Clover, Timothy, Bromegrass, Fescue	16 to 64 fl. oz./acre	Apply 1 - 3 times during active growth.
Root and Tuber Vegetables (Potatoes, Carrots, Onions)	16 to 32 fl. oz./acre	Apply 1 - 3 times during active growth.

Calibrate sprayer to correct rate. If mixing with crop protection products, do a jar test first and apply within 12 hours of mixing. Compatible with liquid fertilizer up to 2 days. Agitate if not used right away. Foliar applications should be made when plants are actively growing.

SHAKE WELL BEFORE USING

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with the directions contained on this label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to, damage to plants, crops, and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops, and personal injury.